



VISION ACADEMY

people | research | education

The association between diabetic retinopathy and other complications of diabetes

Dr. Marc Evans

University Hospital of Llandough, Wales, UK



Financial disclosures

- Honoraria / research support from:
 - Bayer
 - BI
 - Lilly
 - MSD
 - Novartis
 - Novo Nordisk
 - Sanofi
 - Takeda

Vision Academy

Diabetes is associated with a variety of complications

- Diabetes and its complications account for more than 2 million deaths every year
- These impose substantial economic costs on:
 - Patients and their families
 - Healthcare systems
 - National economies (e.g. due to loss of work or wages)
- Together with preventing and delaying the onset of diabetes, early detection and management of complications are important for reducing the health and economic burden of diabetes

Could a greater awareness of diabetic eye disease enable the early detection and management of other diabetic complications?



VISION ACADEMY


people | research | education

The role of the Vision Academy



The role of the Vision Academy

- Currently, there is a lack of communication between the various healthcare professionals involved in the treatment of patients with diabetes
- A literature search was performed to examine the relationship between **DR** and other **microvascular** and **macrovascular** complications of diabetes, with the aim of opening up important communication channels between diabetologists and ophthalmologists in order to improve the coordinated care of patients with diabetes





VISION ACADEMY

people | research | education

DR and microvascular complications of diabetes





Nephropathy: Literature review findings

DR increases the likelihood of having or developing nephropathy¹⁻³

- DR is a significant independent predictor for progression to **micro-** or **macro-albuminuria**¹
 - However, it is unclear whether albuminuria predicts development of DR⁴⁻⁶
- DR is also linked with **declining GFR**^{3,7}
 - The severity of DR at baseline can predict the rate of decline in estimated GFR³


Cumulative incidence of nephropathy with and without DR*,¹



*Follow-up in 537 patients with type 1 diabetes and normal urinary albumin levels at baseline. DR, diabetic retinopathy; GFR, glomerular filtration rate.

1. Rossing P *et al. Diabetes Care* 2002; 25 (5): 859–864. 2. Parving H-H *et al. Kidney Int* 2006; 69 (11): 2057–2063. 3. Rossing K *et al. Kidney Int* 2004; 66 (4): 1596–1605. 4. Kotlarsky P *et al. Int Ophthalmol* 2015; 35 (1): 59–66. 5. Romero-Aroca P *et al. Diabetes Res Clin Pract* 2011; 94 (1): 126–132. 6. Hammes H-P *et al. PLoS One* 2015; 10 (7): e0132492. 7. Moriya T *et al. Diabetes Care* 2013; 36 (9): 2803–2809.

Neuropathy: Literature review findings



DR is associated with cardiac autonomic neuropathy (CAN)^{1,2}

- The risk of CAN is increased in the presence of DR¹
- In patients with T2DM, the stage of DR is independently associated with CAN²

DR is associated with diabetic peripheral neuropathy (DPN)³⁻⁶

- DR is a significant risk factor for DPN, which leads to hospitalization more frequently than other complications of diabetes^{3,7}
 - The risk of DPN is greater in patients with proliferative DR than in those with early stage DR⁴

CAN, cardiac autonomic neuropathy; DPN, diabetic peripheral neuropathy; DR, diabetic retinopathy; T2DM, type 2 diabetes mellitus.

1. Voulgari C *et al. J Diabetes Complications* 2011; 25 (3): 159–167. 2. Huang C-C *et al. J Diabetes Res* 2016; 2016: 6090749. 3. Kostev K *et al. Prim Care Diabetes* 2014; 8 (3): 250–255.

4. Lin I-C *et al. Acta Ophthalmol* 2015; 93 (8): 713–718. 5. Dyck PJ *et al. Diabetes Care* 1999; 22 (9): 1479–1486. 6. Abougambou SSI *et al. Diabetes Metab Syndr* 2015; 9 (2): 98–103.

7. Bansal V *et al. Postgrad Med J* 2006; 82 (964): 95–100.




VISION ACADEMY

people | research | education

DR and macrovascular complications of diabetes



Cerebrovascular complications: Literature review findings



DR is a strong predictor of stroke¹⁻⁴


- Even in mild-to-moderate forms, DR was identified as an independent risk factor for stroke^{2,3}
 - Severe DR independently increased the risk of cerebral infarction and cerebral hemorrhage in patients with T1DM⁴
 - A large prospective study in patients with T2DM found DR to be associated only with small artery ischemic stroke⁵

DR, diabetic retinopathy; T1DM, type 1 diabetes mellitus; T2DM, type 2 diabetes mellitus.

1. Petitti DB *et al. Stroke* 1995; 26 (4): 593–596. 2. Cheung N *et al. Stroke* 2007; 38 (2): 398–401. 3. Klein BEK *et al. Arch Intern Med* 2004; 164 (17): 1917–1924.

4. Hägg S *et al. Diabetes Care* 2013; 36 (12): 4140–4146. 5. Hankey GJ *et al. J Neurol Neurosurg Psychiatry* 2013; 84 (3): 281–287.

Cardiovascular complications: Literature review findings



DR is associated with subclinical atherosclerosis¹⁻³

- The presence of DR was independently associated with **carotid intima media thickness** and **carotid plaques** in patients with T2DM³⁻⁵

DR is an independent risk factor for cardiovascular disease⁶⁻⁹

- In patients with T2DM, DR was associated with an increased risk of **CHD-related events/deaths**^{10,11}


DR (number at risk)	Incident CHD events		Fatal CHD events	
	Number (%) of events	Multivariate HR (95% CI)	Number (%) of events	Multivariate HR (95% CI)
Absent (n=1,242)	153 (12.3)	1.0	19 (1.5)	1.0
Present (n=214)	44 (20.6)	1.99 (1.33–3.00)	13 (6.1)	3.00 (1.27–7.09)

Table adapted from Cheung et al. 2007¹¹

CI, confidence interval; CHD, coronary heart disease; DR, diabetic retinopathy; HR, hazard ratio; T2DM, type 2 diabetes mellitus.


- Liu Y et al. *Diab Vasc Dis Res* 2015; 12 (5): 366–372.
- Saif A et al. *Endocr Pract* 2015; 21 (3): 226–230.
- Rema M et al. *Diabetes Care* 2004; 27 (8): 1962–1967.
- Li L-X et al. *J Diabetes Complications* 2014; 28 (3): 378–385.
- Alonso N et al. *Cardiovasc Diabetol* 2015; 14: 33.
- Gimeno-Orna JA et al. *Am J Cardiol* 2009; 103 (10): 1364–1367.
- Roy MS et al. *Diabet Med* 2007; 24 (12): 1361–1368.
- Targher G et al. *Diabet Med* 2008; 25 (1): 45–50.
- Park G-M et al. *Am J Cardiol* 2014; 113 (5): 765–771.
- Kawasaki R et al. *Ophthalmology* 2013; 120 (3): 574–582.
- Cheung N et al. *Diabetes Care* 2007; 30 (7): 1742–1746.

Peripheral vascular complications: Literature review findings



DR is associated with peripheral arterial disease (PAD)^{1–3}

- Patients with proliferative DR are more likely to have an **abnormal ABPI** than patients with non-proliferative DR³



DR is associated with soft tissue complications^{4–7}

- DR is an independent risk factor for **foot ulceration**, and a key risk factor for **lower extremity amputation**^{4–7}
 - Similarly, patients who have non-healing foot ulcers or who have had lower extremity amputations are high risk of developing and progressing DR^{8,9}

ABPI, ankle-brachial pressure index; DR, diabetic retinopathy; PAD, peripheral arterial disease.

1. Li X *et al. J Diabetes* 2012; 4 (2): 140–146. 2. Chen S-C *et al. PLoS One* 2015; 10 (7): e0134718. 3. Chen Y-W *et al. PLoS One* 2015; 10 (3): e0122022.

4. Baba M *et al. Diabetes Res Clin Pract* 2014; 106 (1): 42–49. 5. Leymarie F *et al. Diabetes Metab* 2005; 31 (6): 603–605. 6. Bruun C *et al. Diabet Med* 2013; 30 (8): 964–972.

7. Parisi MCR *et al. Diabetol Metab Syndr* 2016; 8: 25. 8. Nwanyanwu KH *et al. Diabetes Care* 2013; 36 (6): 1562–1568. 9. Mohammedi K *et al. Cardiovasc Diabetol* 2016; 15 (1): 129.

Session summary

- A literature review of **81 studies** in **>400,000 patients** found that DR can be a predictor of numerous microvascular and macrovascular complications of diabetes



- Conversely, some systemic complications of diabetes are linked to a higher risk of developing or progressing DR



- These findings demonstrate the need for **prompt referral** and **screening** for DR and other complications of diabetes, and highlight the need for a **coordinated approach** to diabetes management

